

August 28, 2000

Does the KARMEN time anomaly originate from a beam-correlated background?

F. Atchison, M. Daum*, P.-R. Kettle, C. Wigger

PSI, Paul-Scherrer-Institut, CH-5232 Villigen-PSI, Switzerland.

*E-mail address: Manfred.Daum@psi.ch

Tel.: +41 56 310 36 68; Fax: +41 56 310 32 94.

ABSTRACT

The time anomaly of neutrino events observed in the KARMEN detector at the ISIS facility is discussed. We show that beam-correlated neutron events are not completely suppressed by the KARMEN lower energy cut and could cause the observed deviation of the measured time distribution from an exponential curve with $\tau = 2.2 \mu s$.

PACS: 13.20.Cz, 14.60.Pq, 14.80.-j

The KARMEN collaboration investigates reactions induced by neutrinos produced in the pulsed neutron spallation source of the ISIS facility at the Rutherford Appleton Laboratory. In 1995, they reported [1] an anomaly in the time distribution of their events clustered around $3.6\,\mu\text{s}$ after beam-on-target with visible energy between 11 and 35 MeV. Further evidence of events exceeding the expected exponential distribution characterized by the muon lifetime has since been reported [2]. The speculative explanation was that these events could be generated by hitherto unobserved neutral massive X -particles originating from the ISIS target via the rare pion decay process,

$$\pi^+ \rightarrow \mu^+ X. \tag{1}$$

From the kinematics of the two-body decay of a pion at rest, and the known masses of the pion and the muon [3], the mass of X was calculated to be 33.9 MeV using the time-of-flight information and the mean flight-path of 17.5 m from the ISIS target to the KARMEN detector. Since the detected energy of the anomalous events is found to be much greater than the kinetic energy of the X -particle, these events are postulated to originate from X -particles which decay with visible energy within the detector via the reaction $X \rightarrow e^+e^-\nu$. The KARMEN collaboration estimated the branching fraction for the decay mode (1) as a function of the X -lifetime, with values of the branching fraction ranging over many decades down to $\sim 10^{-16}$ [1].

Soon after the initial KARMEN publication [1], various hypotheses were put forward as to the nature of the X -particle [4-8], and several searches

undertaken to look for the rare decay (1) [9–12], all with negative results. More recently, additional experiments [13–15] have set even more stringent limits for the branching fraction of the decay mode (1).

In this letter we present arguments for the existence of a beam-correlated neutron background in the KARMEN detector that could contribute to an excess of events¹. Our basis is that the KARMEN detector can respond to the 10.8 MeV γ -energy from neutron capture by ^{14}N . Thermal neutrons will originate from moderation of fast neutrons ($T_n < 10$ MeV) in the liquid scintillator. These fast neutrons are produced from medium energy interactions ($T_n > 100$ MeV) in the KARMEN iron shield.

The liquid scintillator (PPP) used in the KARMEN detector consists of 75 % (volume) paraffin oil, 25 % pseudocumene and 2 g/l PMP (1-phenyl-3-mesityl-2-pyrazoline) [16, 17]. The chemical formulae of the components are (i) paraffin oil, taken as $\text{C}_{11}\text{H}_{24}$, (ii) pseudocumene C_9H_{12} , (iii) PMP $\text{C}_{18}\text{H}_{20}\text{N}_2$. The nuclear densities in the liquid scintillator material (in units of nuclei/ $\text{\AA}^3$) are 0.071 for hydrogen, 0.036 for carbon and $9.1 \cdot 10^{-6}$ for nitrogen leading to the ratio of atomic abundances of

$$n_H : n_C : n_N \approx 7800 : 4000 : 1. \quad (2)$$

It is this nitrogen that can capture neutrons via the reaction

$$^{14}\text{N}(n, \gamma)^{15}\text{N} \quad (3)$$

with an integral γ -energy of 10.8 MeV. The thermal neutron cross-section for

¹The number of excess events reported in Ref. [1] is 83 ± 28 .

this reaction is $\sigma \approx 75 \text{ mb}$. The lower energy cut of events in the KARMEN analysis is 11 MeV [1] with an energy resolution of $\sigma_E/E = 11.5\%/\sqrt{E}$ [16] (E in MeV), which is 3.5% or 380 keV for $E = 11 \text{ MeV}$. Thus, the 10.8 MeV γ -events from reaction (3) will be accepted with an efficiency of about 30 %; we note that in the energy distribution of the excess events [18], about 40 % of the anomalous events, which are selected from an energy window between 11 and 35 MeV, are clustered at about 11 MeV. If such a beam-correlated neutron background is present, the subtraction of purely cosmic background events is not sufficient.

The data fit published by the KARMEN collaboration gives a value for the time constant $\tau_\mu = (2.62 \pm 0.18) \mu\text{s}$ [1] which differs by more than two standard deviations from the expected value of the muon lifetime ($2.19703 \pm 0.00004 \mu\text{s}$). We have refitted the KARMEN data of Ref. [1] with an additional background of the form:

$$y(t) = A \cdot e^{-t/\tau_\mu} + B, \quad (4)$$

and fixing τ_μ to the value of $2.2 \mu\text{s}$. The result of our fit is shown in Fig. 1. This fit has a χ^2 of 16.4 for 17 DOF and the values of the two free parameters found are $A = (271 \pm 16) \text{ events}/0.5\mu\text{s}$, $B = (7.2 \pm 3.0) \text{ events}/0.5\mu\text{s}$. The value for B , greater than zero by 2.4 standard deviations, can be interpreted as an indication for the presence of an additional background with $137 \pm 57 \text{ events}^2$.

²A similar fit was made to the combined data from KARMEN1 and KARMEN2 [18].

From the thermal capture cross-sections of hydrogen (333 mb), carbon (3.5 mb) and nitrogen (75 mb for n,γ and 1810 mb for n,p) and using the ratio of atomic abundances (2), we deduce the capture cross-section of the scintillator to be about 220 mb and the probability for reaction (3) to be $P \approx 3 \cdot 10^{-5}$. This means that $1/P \approx 3 \cdot 10^4$ thermal neutrons would be needed to create photons of reaction (3) with 10.8 MeV of deposited visible energy. In order to pass the 11 MeV cut in the KARMEN analysis, this number has to be a factor of 3 larger, i.e. about $1 \cdot 10^5$.

Because of the neutron absorbers around each scintillator element (gadolinium and borated polyethylene) [1], thermal neutrons cannot enter the KARMEN detector and hence have to be produced by moderating fast neutrons in the scintillator fluid itself. Since the hydrogen density in the PPP is similar to that for water, the moderation characteristics should be alike. ‘Age theory’ [19] predicts an average time for thermalization of a few microseconds. The time variation of the background signal will be complicated by the $1/v$ -tailback of the capture cross-section. Thus, the contributions to the background signal, will start well before thermalization is complete.

The events of Fig. 1 originate from 6560 C, i.e. $4 \cdot 10^{22}$ protons hitting the ISIS target [1]. Each proton produces about 0.05 medium energy neutrons per steradian at 90 degree production angle [20]. The KARMEN detector covers $5 \cdot 10^{-3}$ of the solid angle. The material between the target and the KARMEN detector provides about $4600 \text{ g}\cdot\text{cm}^{-2}$ of shielding [1,16,20,21], and the shielding length for medium energy (above $\sim 100 \text{ MeV}$) neutrons is in the

range 120 to 160 $\text{g}\cdot\text{cm}^{-2}$ [22]. This leads to the estimate that between $5 \cdot 10^7$ and $3 \cdot 10^3$ medium energy neutrons enter the KARMEN detector. Medium energy neutrons contribute 0.4 % of the total neutron flux escaping from an iron shield [20], and 98 % of all neutrons have an energy below 10 MeV. With these numbers, we get the contributions to our probability estimate of a beam-correlated neutron background to be detected in the KARMEN time window ($10\mu\text{s}$) listed in Table 1.

From this table, we calculate the suppression factor of beam-correlated neutrons to be between $1 \cdot 10^{-24}$ and $2 \cdot 10^{-20}$ for the two extremes of the shielding length. Thus, for a total number of protons ($4 \cdot 10^{22}$), the corresponding expected number of neutron events originating from a beam-correlated background is between 0.06 and 800 which brackets the number of background events (137 ± 57).

For a more refined background estimate, one has to consider that in a pulsed neutron spallation source, a beam-correlated neutron background has a time structure originating from the duty cycle of the primary proton beam and the kinetic energy of the neutrons on their way through the shielding. The neutron capture rate of gadolinium absorber dissolved in water was measured as a function of time after injection of a fast neutron pulse [23] and corresponds to a linear rise from zero to about $6\mu\text{s}$ followed by a constant rate at longer times³. The time variation of the neutron capture rate in the liquid

³In fact, an exponential fall with a time constant of $\sim 100\mu\text{s}$ [16] (the thermal neutron lifetime in the KARMEN scintillator) is to be expected.

scintillator is expected to be similar. This behavior has been approximated by the function

$$y(t) = A \cdot e^{(-t/2.2\mu s)} + B \cdot t/\tau_c \quad (5)$$

for $t < \tau_c$ and

$$y(t) = A \cdot e^{(-t/2.2\mu s)} + B \quad (6)$$

for $t > \tau_c$. Here, A is the initial rate of the exponential decay, B is the beam-correlated background from reaction (3), and τ_c is the linear risetime of the neutron capture rate. A fit to the data of Ref. [1] results in a χ^2 of 15.5 for 16 DOF. The fitted values of the free parameters were $A = (277 \pm 14)$ events/ $0.5\mu s$, $B = (7.3 \pm 2.8)$ events/ $0.5\mu s$, and $\tau_c = (3.4 \pm 0.5)\mu s$. This fit, shown in Fig. 2, leads to 120 ± 42 neutron events from a beam-correlated background. Although the statistical significance is similar to the previous fit, it gives a possible explanation for an enhancement in the spectrum due to the neutron capture rate in the scintillator reaching its maximum some 3 to $4\mu s$ after beam on target.

After the upgrade of the KARMEN shielding (1996/97), any beam-correlated background from neutrons should be further suppressed due to additional polyethylene, veto scintillation counters, and iron shielding. Thus, the corresponding ‘time anomaly’ in the KARMEN data [1, 18] before 1997, should be less significant in the data after 1997, as was presented in Ref. [24].

In conclusion, it can at present not be ruled out that the number of KARMEN excess events and their time distribution according to Ref. [1] are

consistent with the hypothesis of originating from a beam-correlated neutron background involving neutron capture in nitrogen via reaction (3). A more elaborate analysis of the time structure of such a neutron background [2, 18] is beyond the scope of this letter, however, we would like to mention the existence of deep minima in the total cross section of iron, e.g. around 81 keV [25]. At this energy, neutrons travel through iron with a mean free path of several meters with a velocity of $4\text{ m}/\mu\text{s}$ and have a TOF of $\sim 2\mu\text{s}$ from the surface of the ISIS biological shielding to the KARMEN detector. These neutrons have a time structure similar to that of the proton beam on the ISIS target and so could lead to a localized enhancement of the beam-correlated background event-rate.

The authors thank G. H. Eaton, R. Eichler, R. Frosch, and V. Markushin for valuable discussions and interest. This work was supported by the Schweizerischer Nationalfonds.

References

- [1] KARMEN collaboration, B. Armbruster et al., Phys. Lett. B 348, 19 (1995).
- [2] KARMEN collaboration, B. Zeitnitz, Proceedings of the 17th International Workshop on Weak Interactions and Neutrinos, WIN'99, Cape Town, ZAR, 150 (1999).
- [3] Review of Particle Properties, Eur. Phys. Jour. C 3 (1998).
- [4] V. Barger et al., Phys. Lett. B 352, 365 (1995);
V. Barger et al., Phys. Lett. B 356, 617(E) (1995).
- [5] M. M. Guzzo et al., Phys. Rev. D 53, 2851 (1996).
- [6] D. Choudhury and S. Sarkar, Phys. Lett. B 374, 87 (1996).
- [7] S. N. Gninenko and N. V. Krasnikov, Phys. Lett. B 434, 163 (1998).
- [8] J. Govaerts et al., Phys. Lett. B 389, 700 (1996).
- [9] M. Daum et al., Phys. Lett. B 361, 179 (1995).
- [10] R. Bilger et al., Phys. Lett. B 363, 41 (1995).
- [11] D. A. Bryman and T. Numao, Phys. Rev. D 53, 558 (1996).
- [12] R. Bilger et al., Phys. Lett. B 446, 363 (1999).
- [13] J. A. Formaggio et al., Phys. Rev. Lett. 84, 4043 (2000).

- [14] M. Daum et al., Phys. Rev. Lett., (2000), in print,
see also hep-ex/0008014.
- [15] NOMAD collaboration, Poster at ν 2000, Sudbury, Canada (2000).
- [16] G. Drexlin et al., Nucl. Instr. and Meth. in Phys. Res. A289, 490 (1990).
- [17] H. Güsten et al., J. Phys. Chem. 82, 459 (1978).
- [18] Ch. Oehler, PhD-thesis, University Karlsruhe, 118 (1999).
see also http://www-ik1.fzk.de/www/karmen/publist_e.html
- [19] K. H. Beckurts, K. Wirtz, “Neutron Physics”, Springer Verlag OHG
Berlin, 145ff, (1964).
- [20] F. Atchison, Rutherford Appleton Laboratory Report RL-81-006 (1981).
- [21] B. Boardman, Rutherford Appleton Laboratory Report RL-82-006
(1982).
- [22] IAEA Technical Reports Series No. 283, 280ff (1988).
- [23] K. H. Beckurts, K. Wirtz, “Neutron Physics”, Springer Verlag OHG
Berlin, 401ff, (1964).
- [24] KARMEN collaboration, K. Eitel, ν 2000, Sudbury, Canada (2000),
see also <http://www-ik1.fzk.de/www/karmen/karmen.html>.

[25] G. Rohr et al, Nuclear Data for Reactors Conf., Paris, 137 (1966).

see also D. I. Garber and R. R. Kinsey, Neutron Cross Sections, Volume II, Curves, Brookhaven National Laboratory, 138ff (1976).

Figure Captions

Figure 1:

KARMEN data of Ref. [1] fitted with the function $y(t) = A \cdot e^{-t/2.2\mu s} + B$. The smooth line represents the fitting function. The dashed line is an exponential with $\tau_\mu = 2.2\mu s$, and the dotted line represents an additional background. The values of the fitted parameters are $A = (271 \pm 16) \text{ events}/0.5\mu s$, $B = (7.2 \pm 3.0) \text{ events}/0.5\mu s$; the χ^2 of the fit is 16.4 for 17 DOF.

Figure 2:

KARMEN data of Ref. [1] fitted with the function $y(t) = A \cdot e^{-t/2.2\mu s} + B \cdot t/\tau_c$ for $t \leq \tau_c$ and $y(t) = A \cdot e^{-t/2.2\mu s} + B$ for $t > \tau_c$. The smooth line represents the fitting function. The dashed line is an exponential with $\tau_\mu = 2.2\mu s$, and the dotted line represents the time dependence of our estimate of neutron capture in the KARMEN scintillator from reaction (3). The values of the fitted parameters are $A = (277 \pm 14) \text{ events}/\mu s$, $B = (7.3 \pm 2.8) \text{ events}/\mu s$, and $\tau_c = (3.4 \pm 0.5) \mu s$; the χ^2 of the fit is 15.5 for 16 DOF.

Figure 1:




Figure 2:




Table 1: Factors used for estimating the beam-correlated neutron background

Number of medium energy neutrons directed towards KARMEN	0.05/sr/proton
Solid angle for the KARMEN detector	0.0628 sr
Shielding factors for 4600 g/cm ² and shielding lengths λ of 120 g/cm ² – 160 g/cm ²	$2 \cdot 10^{-17} - 3 \cdot 10^{-13}$
Fraction of neutrons escaping the KARMEN steel shield in the medium energy region (0.4 %), in the fast neutron region (98 %); the number of fast neutrons per medium energy neutron is	$0.98/0.004 \approx 250$
Shielding effect for fast neutrons of the borated polyethylene layer and the active outer shield	0.25
Fraction of neutrons captured by PPP	0.78
Probability of the capture channel $^{14}\text{N}(n, \gamma)^{15}\text{N}$	$3 \cdot 10^{-5}$
Detection efficiency for 10.8 MeV (energy cut 11 MeV)	0.3
Fraction of captures within the KARMEN time window	0.05